arXiv:1404.7420v4 [math-ph] 11 Jul 2014

Master symmetry and time dependent symmetries of the differential-difference KP equation

Farbod Khanizadeh
School of Mathematics, Statistics & Actuarial Science
University of Kent, Canterbury, UK

Abstract

We first obtain the master symmetry of the differential-difference KP equation.Then we show how this master symmetry, through $sl(2,\mathbb{C})$-representation of the equation, can construct generators of time dependent symmetries.

1 Introduction

The differential-difference KP (DDKP) equation is a lattice field equation of the following form [1, 2, 3, 4]:

$$u_t = u_{xx} + 2uu_x + 2(\mathcal{S}-1)^{-1}u_{xx} = K\,, \tag{1.1}$$

where the dependent variable $u := u(n,x,t)$ is a smooth function of continuous varibles x, t and discrete variable $n \in \mathbb{Z}$. We use the notation $u_{m,j} =: D_x^j u(n+m,t,x) = \mathcal{S}^m D_x^j u(n,t,x)$ where $\mathcal{S}$ is the linear shift operator and D_x denotes the derivative with respect to x. The DDKP equation is also appeared in [5, 6]. In [5] it is constructed through the procedure of central extension and the author refers to the equation as the lattice field Benjamin-Ono. For the links between hierarchy of equation (1.1) and the continuous Kadomtsev-Petviashvili hierarchy one can read [7].

Notice that equation (1.1) involves $(\mathcal{S}-1)^{-1}$ and is no longer local. The nonlocality is the main obstacle to directly extend the theories of (1+1)-dimensional nonlinear evolutionary equations to (2+1)-dimensional equations.

Master symmetries are tools by which, through a recursive process, one can construct a hierarchy of infinitely many commuting generalised symmetries and guarantee the integrability. The notion of master symmetry was first introduced in [8] and later studied in [9, 10]. The efficiency of these tools can be clearly seen in the (2+1)-dimensional case for which recursion operators of equations have more complicated structure than in (1+1)-dimension [11, 12].

Master symmetries of (1+1)-dimensional differential and differential-difference equations are not always local. For example the KdV

$$u_t = u_{xxx} + 6uu_x\,,$$

admits the following nonlocal master symmetry[13, 14, 15]:

$$\tau = x(6uu_x + u_{xxx}) + 4u_{xx} + 8u^2 + 2u_x D_x^{-1}u\,. \tag{1.2}$$

Master symmetries are also related closely to time dependent symmetries of evolutinary equations. In [9, 16] this relation is discussed along with examples of continous case. It is shown having a master symmetry of evolution equation, one can produce an infinite number of time dependent symmetries which are polynomial in time.

In this paper our focus is on the DDKP equation. We write the equation in the form in which a quasi-local term appears. Then we introduce briefly the concept of master symmetries and derive the master symmetry of the equation. The paper ends with the application of the master symmetry in constructing generators of the time dependent symmetries of the DDKP equation. This relation is illustrated through the $\mathrm{sl}(2,\mathbb{C})$-representation of the DDKP equation.

2 The differential-difference KP equation and its master symmetry

We can rewrite the differential-difference KP (DDKP) equation (1.1) in the following form:

$$u_t = u_{xx} + 2uu_x + 2\Theta(u_x) = K\,, \tag{2.1}$$

where $\Theta := (\mathcal{S}-1)^{-1}D_x$ and $\Theta(u_x)$ is called a quasi-local function.

The concept of quasi-local functions was first introduced for the continous case in 1998 by Mikhailov and Yamilov [17].They noticed that the operators D_x^{-1} and D_y^{-1} never appear alone but always in pairs like $D_x^{-1}D_y$ and $D_y^{-1}D_x$ for all known integrable equations and their hierarchies of symmetries and conservation laws. In [18], using the symbolic representation, the author proved this observation is true for certain type of integrable equations. Here we denote $\hat{\mathfrak{G}}$ for the space of quasi-local polynomials which depend on $u_{i,j}$ and contain the operator Θ. We also define $\mathfrak{G}$ as the extension of $\hat{\mathfrak{G}}$ to the space with coefficients from the set $\{n, x\}$. For more details and discussion on the structure of $\hat{\mathfrak{G}}$ and $\mathfrak{G}$ one can read [19].

For two elements F and G in $\mathfrak{G}$ we define the Lie bracket

$$[F,G] := F_*(G) - G_*(F)\,, \tag{2.2}$$

where F_* is called the Fréchet derivative of F and is defined by:

$$F_* = \sum_{i,j} \frac{\partial F}{\partial u_{i,j}} \mathcal{S}^i D_x^j\,. \tag{2.3}$$

The anti-symmetric axiom is clear and bilinearity is deduced from the linearity of derivations and the shift operator. One can also prove the Jacobi identity through a direct calculation.

Definition 1. *We call an element $G \in \hat{\mathfrak{G}}$ a generalised symmetry of equation (2.1) if and only if $[K,G] = 0$. Furthermore equation (2.1) is called integrable if and only if it possesses infinitely many commuting generalised symmetries.*

As mentioned in the introduction, the master symmetry is a quantity by which one can construct a hierarchy of commuting symmetries. For a comprehensive description of master symmetries one can read [8, 9, 20, 21, 22, 23].

Definition 2. *An element $W \in \mathfrak{G}$ is called a master symmetry of equation (2.1) if and only if*

$$[K,[K,W]] = 0, \qquad [W,K] \in \hat{\mathfrak{G}}\,. \tag{2.4}$$

In practice we are interested to find a W such that its recursive action on K is well-defined and produces a new element in $\hat{\mathfrak{G}}$. In the following theorem we provide the explicit form of the master symmetry of equation (2.1).

Theorem 1. *The element $W \in \mathfrak{G}$ of the form*

$$W = xu_{xx} + 2xuu_x + 2x\Theta(u_x) + nu_x + u^2 + 3\Theta(u)\,, \tag{2.5}$$

is the master symmetry of the DDKP equation (2.1). Furthermore the recursive action of W on the equation produces a commutative space of symmetries.

Proof. According to Definition 2, to say W is a master symmetry we need to show $[K,[W,K]] = 0$ where $[W,K]$ lies in the space $\hat{\mathfrak{G}}$. This implies that $[W,K]$ should be a generalised symmetry of equation (2.1). The first non-trivial symmetry of (2.1) is given in [5]. We rewrite it in our notation as follows:

$$\begin{aligned} G &= u_{xxx} + 3uu_{xx} + 3u_x^2 + 3u^2u_x + 3\Theta(uu_x) + 3u_x\Theta(u) \\ &+ 3u\Theta(u_x) + 3\Theta^2(u_x) + 3\Theta(u_{xx})\,. \end{aligned} \tag{2.6}$$

From (2.3) one can check that the bracket $[K,G]$ vanishes and $[W,K] = -2G$, which proves the first part.

The second part of the theorem is thoroughly discussed in [19]. There, the author refers to the theorem namely τ-scheme by I. Dorfman [24] and shows how W meets the conditions of the τ-scheme. □

In the next section we give the description of an irreducible $sl(2,C)$-representation by which we can construct the time dependent symmetries of the DDKP equation.

3 Time dependent symmetries

In [16] it was shown how the representation of Burgers, Ibragimov-Shabat and KP equation can help one to construct the time dependent symmetries. In what follows, using the Lie algebra $sl(2,\mathbb{C})$ we present such a construction for equation (2.1). We call $G(u,t)$ a time dependent symmetry of (2.1) if and only if

$$\frac{\partial G}{\partial t} = [G,K]\,. \tag{3.1}$$

We find an appropriate element $G_0 \in \hat{\mathfrak{G}}$ such that for some ℓ we get $ad_K^\ell(G_0) = 0$, where

$$ad_K^\ell G_0 := [\underbrace{K\cdots,[K,[K}_{\ell-times},G_0]]]\,,$$

Then it can be shown $G(u,t) = \exp(-tad_K)(G_0)$ is a time dependent symmetry of equation (2.1) [9, 16]. To begin with, we consider

$$M = K, \quad N = -\frac{1}{2}x, \quad H = -xu_x - u\,. \tag{3.2}$$

Calculating the Lie brackets yields

$$[M,N] = H, \quad [H,N] = -2N, \quad [H,M] = 2M\,. \tag{3.3}$$

Therefore we have the following Lie algebra isomorphisms:

$$sl(2,C) \cong span\{M,N,H\}$$

We know from the representation theory of $sl(2,\mathbb{C})$ (more details can be found in [25, 26]) that if V is a finite dimensional irreducible representation, then there is a vector $\upsilon_0 \in V$ where $H\upsilon_0 = \lambda\upsilon_0$ and $M\upsilon_0 = 0$ and furthermore V is spanned by the $\lambda+1$ linear independent vectors $\{\upsilon_0, ad_N\upsilon_0, ad_N^2\upsilon_0, \cdots, ad_N^\lambda\upsilon_0\}$. The element υ_0 is called the highest weight vector.

Since $[M,K]=0$ and $[H,K]=2K$, by setting $\upsilon_0 = K$ we obtain a three dimensional representation of $sl(2,\mathbb{C})$ spanned by $V_3 = span\{K, ad_N K, ad_N^2 K\}$. Now let N_2 be the master symmetry of the DDKP equation. Hence

$$[M, ad_{N_2}^m K] = 0\,.$$

The following theorem provides the basis of $sl(2,\mathbb{C})$-representations with different dimension. For the proof one can read [16].

Theorem 2. *For elements defined in (3.2) the following relation holds:*

$$[H, ad_{N_2}^m K] = (m+2) ad_{N_2}^m K\,.$$

This theorem implies that for different value of m, $ad_{N_2}^m K$ is a highest weight vector of the $(m+3)$-dimensional representation

$$V_{m+3} = span\{ad_{N_2}^m K, ad_N ad_{N_2}^m K, ad_N^2 ad_{N_2}^m K, \cdots, ad_N^{m+2} ad_{N_2}^m K\}\,,$$

where

$$ad_N^{m_2} ad_{N_2}^{m_1} K = [\underbrace{N,[N,\cdots,[N}_{m_2-times}, \overbrace{[N_2,[N_2,\cdots[N_2}^{m_1-times}, K]]]]]]\,.$$

Therefore starting from u_x we can draw the following diagram for the DDKP equation in which the first element of each row is the symmetry of equation (2.1) and the horizontal lines contain the representations of $sl(2,\mathbb{C})$:

$$\begin{array}{ccccccccc}
u_x & \xrightarrow{N} & ad_N u_x & & & & & & \\
\downarrow N_2 & & & & & & & & \\
K & \xrightarrow{N} & ad_N K & \xrightarrow{N} & ad_N^2 K & & & & \\
\downarrow N_2 & & & & & & & & \\
ad_{N_2} K & \xrightarrow{N} & ad_N ad_{N_2} K & \xrightarrow{N} & ad_N^2 ad_{N_2} K & \xrightarrow{N} & ad_N^3 ad_{N_2} K & & \\
\downarrow N_2 & & & & & & & & \\
ad_{N_2}^2 K & \xrightarrow{N} & ad_N ad_{N_2}^2 K & \xrightarrow{N} & ad_N^2 ad_{N_2}^2 K & \xrightarrow{N} & ad_N^3 ad_{N_2}^2 K & \xrightarrow{N} & ad_N^4 ad_{N_2}^2 K \\
\downarrow N_2 & & & & & & & & \\
\vdots & & & & & & & &
\end{array}$$

In the following theorem we see that horizontal lines are not just elements of the representation but generators for time dependent symmetries of equation (2.1).

Theorem 3. *Consider $ad^m_{N_2}K \in V_{m+3}$, then the basis elements*

$$ad^\ell_N ad^m_{N_2}K\,,$$

commute with $K^{\ell+1}, (0 \leq \ell \leq m+2)$. In other words we have

$$ad^{\ell+1}_K(ad^\ell_N ad^m_{N_2}K) = 0\,.$$

Proof. The proof is by induction on ℓ. For $\ell = 0$ as all $ad^m_{N_2}K$ are symmetries of (2.1), we have

$$[K, ad^m_{N_2}K] = 0\,.$$

Assume the statement holds for

$$ad^{\ell+1}_K(ad^\ell_N ad^m_{N_2}K) = 0\,,$$

then we have

$$\begin{aligned} ad^{\ell+2}_K ad^{\ell+1}_N ad^m_N K &= [K, ad^{\ell+1}_K[N, ad^\ell_N ad^m_{N_2}K]] \\ &= [K, [ad^{\ell+1}_K N, ad^{\ell+1}_K ad^\ell_N ad^m_{N_2}K]] = 0\,. \end{aligned}$$

□

Hence each basis element, lying on the horizontal line, provides a generator for the time dependent symmetries of the DDKP equation. Bellow we give two examples of such symmetries

$$\begin{aligned} &\exp(-tad_K)(ad_N K) = [K,N] - t[K,[K,N]] = (-xu_x - u) + 2t(u_{xx} + 2uu_x + 2\Theta(u_x))\,, \\ &\exp(-tad_K)(ad^2_N K) = ad^2_N K - t(ad_K ad^2_N K) + \frac{t^2}{2!}(ad^2_K ad^2_N K) = x - 2t(xu_x - u) \\ &+4t^2(u_{xx} + 2uu_x + 2\Theta(u_x))\,. \end{aligned}$$

4 Conclusion

In this paper we showed how the master symmetry of the DDKP equation can produce an infinite hierarchy of commuting symmetries. We also demonstrated how systematically, using the Lie algebra $sl(2,\mathbb{C})$, one can obtain time dependent symmetries that are polynomial in t. The Lie algebra $sl(2,\mathbb{C})$ is constructed around the scaling symmetry and the master symmetry of a given equation. This structure can be also used for (1+1)-dimensional equations in which the equation, symmetries and the basis elements of $sl(2,\mathbb{C})$ are all local.

Acknowledgements

The author would like to express his sincere appreciation to his supervisor Dr J. P. Wang for her guidance and advice throughout this paper. I would like to thank Dr. G. Papamikos for many helpful discussions and comments. This work was also carried out during my PhD study in SMSAS at the University of Kent and I am grateful for supports received from the School.

References

[1] F.W. Nijhoff, H.W. Capel, G.L. Wiersma and G.R.W. Quispel. Bäcklund transformations and three-dimensional lattice equations. *Physics Letters A*, 105(6):267–272, 1984.

[2] E. Date, M. Jimbo and T. Miwa. Method for generating discrete soliton equations.I. *Journal of the Physical Society of Japan*, 51:4116–4124, 1982.

[3] F.W. Nijhoff. The direct linearizing transform for three-dimensional lattice equations. *Physica D: Nonlinear Phenomena*, 18(1):380–381, 1986.

[4] T. Tamizhmani, V.S. Kanaga and K.M. Tamizhmani. Wronskian and rational solutions of the differential-difference KP equation. *Journal of Physics A: Mathematical and General*, 31(37):7627–7633, 1998.

[5] M. Blaszak and A. Szum. Lie algebraic approach to the construction of (2+1)– dimensional lattice–field and field integrable Hamiltonian equations. *Journal of Mathematical Physics*, 42(1):225–259, 2001.

[6] S. Kanaga and K.M. Tamizhmani. Lax pairs, symmetries and conservation laws of a differential-difference equation–Sato's approach. *Chaos, Solitons & Fractals*, 8(6):917–931, 1997.

[7] W. Fu, L. Huang, K.M. Tamizhmani, and D. Zhang. Integrability properties of the differential-difference Kadomtsev–Petviashvili hierarchy and continuum limits. *Nonlinearity*, 26(12):3197, 2013.

[8] A.S. Fokas and B. Fuchssteiner. The hierarchy of the Benjamin– Ono equation. *Physics Letters A*, 86(6-7):341–345, 1981.

[9] B. Fuchssteiner. Mastersymmetries, higher order time-dependent symmetries and conserved densities of nonlinear evolution equations. *Progress of Theoretical Physics*, 70(6):1508–1522, 1983.

[10] W. Oevel and B. Fuchssteiner. Explicit formulas for symmetries and conservation laws of the Kadomtsev– Petviashvili equation. *Phys. Lett. A*, 88(7):323–327, 1982.

[11] A.S. Fokas and P.M. Santini. Recursion operators and Bi-hamiltonian structures in multidimensions .II. *Communications in Mathematical Physics*, 116(3):449–474, 1988.

[12] P.M. Santini and A.S. Fokas. Recursion operators and Bi-hamiltonian structures in multidimensions . I. *Communications in Mathematical Physics*, 115(3):375–419, 1988.

[13] I.Ya. Dorfman. Dirac structures of integrable evolution equations. *Physics Letters A*, 125(5):240–246, 1987.

[14] A.S. Fokas. Symmetries and integrability. *Studies in Applied Mathematics*, 77:253–299, 1987.

[15] W. Oevel. *Rekursionsmechanismen für Symmetrien und Erhaltungssätze in integrablen Systemen*. PhD thesis, Universität–Gesamthochschule Paderborn, Paderborn, 1984.

[16] J.A. Sanders and J.P. Wang. On integrability of evolution equations and representation theory. *Contemporary Mathematics*, 285:85–99, 2001.

[17] A.V. Mikhailov and R.I. Yamilov. Towards classification of $(2+1)$-dimensional integrable equations. Integrability conditions. I. *Journal of Physics. A. Mathematical and General*, 31(31):6707–6715, 1998.

[18] J.P. Wang. On the structure of $(2+1)$–dimensional commutative and noncommutative integrable equations. *Journal of Mathematical Physics*, 47(11):113508, 2006.

[19] F. Khanizadeh. *Symmetry structure for differential–difference equations*. PhD thesis, University of Kent, Canterbury, 2013.

[20] B. Fuchssteiner. Mastersymmetries for completely integrable systems in statistical mechanics. In *Applications of Field Theory to Statistical Mechanics*, pages 305–315. Springer, 1985.

[21] B. Fuchssteiner and W.X. Ma. An approach to master symmetries of lattice equations. In *Symmetries and integrability of difference equations (Canterbury, 1996)*, pages 247–260. Cambridge Univ. Press, Cambridge, 1999.

[22] W. Oevel and M. Falck. Master symmetries for finite dimensional integrable systems. *Progress of theoretical physics*, 75(6):1328–1341, 1986.

[23] W. Oevel and B. Fuchssteiner. Explicit formulas for symmetries and conservation laws of the Kadomtsev–Petviashvili equation. *Physics Letters A*, 88(7):323–327, 1982.

[24] I.Ya. Dorfman. *Dirac structures and integrability of nonlinear evolution equations*. John Wiley & Sons Ltd., Chichester, 1993.

[25] K. Erdmann and M.J. Wilson. *Introduction to Lie Algebras*. Undergraduate Mathematics Series. Springer, Verlag London, 2006.

[26] J.E. Humphreys. *Introduction to Lie algebras and representation theory*. Springer New York, 1972.